

FIELD ALGEBRAS

BOJKO BAKALOV AND VICTOR G. KAC

Dedicated to Ernest Borisovich Vinberg on the occasion of his 65th birthday.

ABSTRACT. A field algebra is a “non-commutative” generalization of a vertex algebra. In this paper we develop foundations of the theory of field algebras.

0. INTRODUCTION

Roughly speaking, the notion of a vertex algebra [B1] is a generalization of the notion of a unital commutative associative algebra where the multiplication depends on a parameter. (In fact, in [B2] vertex algebras are described as “singular” commutative associative rings in a certain category.) More precisely, an operator of left multiplication on a vertex algebra V by an element $a \in V$ is a *field* $Y(a, z) = \sum_{n \in \mathbb{Z}} a_{(n)} z^{-n-1}$, where $a_{(n)} \in \text{End } V$, and one requires that $Y(a, z)b$ is a Laurent series in z for any two elements $a, b \in V$. The role of a unit element of an algebra is played in this context by a *vacuum vector* $|0\rangle \in V$, satisfying

$$(\text{vacuum axioms}) \quad Y(|0\rangle, z) = I_V, \quad Y(a, z)|0\rangle = e^{zT}a,$$

where $I_V \in \text{End } V$ is the identity operator and $T \in \text{End } V$. A linear map $a \mapsto Y(a, z)$ of a vector space V with a vacuum vector $|0\rangle$ to the space of $\text{End } V$ -valued fields satisfying

$$(\text{translation invariance}) \quad [T, Y(a, z)] = Y(Ta, z) = \partial_z Y(a, z),$$

is called a *state-field correspondence*. This notion is an analogue of a unital algebra.

For example, if V is an ordinary algebra with a unit element $|0\rangle$ and T is a derivation of V , then the formula

$$(0.1) \quad Y(a, z)b = (e^{zT}a)b, \quad a, b \in V$$

defines a state-field correspondence (and it is easy to show that all of them with the property that $Y(a, z)$ is a formal power series in z , are thus obtained).

Furthermore, the associativity property of the algebra V is equivalent to the following property of the state-field correspondence (0.1):

$$(0.2) \quad Y(Y(a, z)b, -w)c = Y(a, z-w)Y(b, -w)c, \quad a, b, c \in V,$$

and the commutativity property of V is equivalent to:

$$(0.3) \quad Y(a, z)b = e^{zT}(Y(b, -z)a), \quad a, b \in V.$$

It turns out that for a general vertex algebra V identity (0.2) holds only after one multiplies both sides by $(z-w)^N$ where N is sufficiently large (depending on

Date: April 23, 2002. Revised May 6, 2002.

The first author was supported by the Miller Institute for Basic Research in Science.

The second author was supported in part by NSF grant DMS-9970007.

a, b, c):

$$(0.4) \quad (z-w)^N Y(Y(a, z)b, -w)c = (z-w)^N Y(a, z-w)Y(b, -w)c, \quad N \gg 0.$$

One of the equivalent definitions of a *vertex algebra* is that it is a state-field correspondence satisfying (0.3) and (0.4) (see Theorem 6.3).

A *field algebra* is a state-field correspondence satisfying only the associativity property (0.4). We believe that this is the right analogue of a unital associative algebra. In the present paper we are making the first steps towards a general theory of field algebras.

The trivial examples of field algebras are provided by state-field correspondences (0.1): this is a field algebra if and only if the underlying algebra V is associative. The simplest examples of non-trivial field algebras are tensor products of field algebras (0.1) with vertex algebras. A special case of this is the algebra of matrices with entries in a vertex algebra. Other examples are provided by a smash product of a vertex algebra and a group of its automorphisms. Thus, many important ring-theoretic constructions with vertex algebras become possible in the framework of field algebras.

One of our main results is the construction of a canonical structure of a field algebra in a tensor algebra $\mathcal{T}(R)$ over a Lie (even Leibniz) conformal algebra R (Theorem 5.1). Imposing relation (0.3) on $\mathcal{T}(R)$ gives the enveloping vertex algebra $\mathcal{U}(R)$ of R (cf. [K, GMS]).

We also establish the field algebra analogues of the density and duality theorems in the representation theory of associative algebras (see Theorems 8.5 and 8.6) and discuss the Zhu algebra construction [Z] in the framework of field algebras.

Note that the “field algebras” considered in [K, Sec. 4.11] are defined by a stronger than associativity axiom; we call them strong field algebras in the present paper. Surprisingly, it turns out that they are almost the same as vertex algebras (see Theorem 7.4), although the “trivial” field algebras (0.1) are automatically strong field algebras. In the present paper the results of [K, Sec. 4.11] on field algebras are corrected.

The first examples of non-trivial field algebras (i.e., different from (0.1)) that are not vertex algebras were constructed in [EK]. The “quantum vertex algebras” of [EK] are field algebras satisfying in addition a certain “braided commutativity” generalizing (0.3). The relation of the present work to the paper [EK], and also to [B2] and [FR], will be discussed in a subsequent paper.

1. STATE-FIELD CORRESPONDENCE

Let V be a vector space (referred to as the *space of states*). Recall that a (End V -valued) *field* is an expression of the form

$$a(z) = \sum_{n \in \mathbb{Z}} a_{(n)} z^{-n-1},$$

where z is an indeterminate, $a_{(n)} \in \text{End } V$, and for each $v \in V$ one has:

$$a_{(n)} v = 0 \text{ for } n \gg 0,$$

i.e., $a(z)v$ is a Laurent series in z .

Denote by $g\ell f(V)$ the space of all End V -valued fields. For each $n \in \mathbb{Z}$ one defines the n -th *product* of fields $a(z)$ and $b(z)$ by the following formula:

$$(1.1) \quad a(z)_{(n)} b(z) = \text{Res}_x (a(x)b(z)i_{x,z}(x-z)^n - b(z)a(x)i_{z,x}(x-z)^n).$$

Here $i_{x,z}$ (respectively $i_{z,x}$) stands for the expansion in the domain $|x| > |z|$ (respectively $|z| > |x|$):

$$i_{x,z}(x-z)^n = \sum_{j=0}^{\infty} \binom{n}{j} x^{n-j} (-z)^j,$$

while

$$i_{z,x}(x-z)^n = \sum_{j=0}^{\infty} \binom{n}{j} x^j (-z)^{n-j}.$$

It is easy to see that the space of fields $glf(V)$ is closed under all n -th products and also under the derivation by the indeterminate.

Recall that

$$(1.2) \quad \delta(x-z) = (i_{x,z} - i_{z,x})(x-z)^{-1} = \sum_{j \in \mathbb{Z}} x^j z^{-j-1}$$

is the formal delta-function, characterized by the property:

$$\text{Res}_x a(x) \delta(x-z) = a(z) \quad \text{for } a(x) \in glf(V).$$

Formula (1.1) is equivalent to the following two formulas for $n \in \mathbb{Z}_+$:

$$\begin{aligned} a(z)_{(n)} b(z) &= \text{Res}_x [a(x), b(z)] (x-z)^n, \\ a(z)_{(-n-1)} b(z) &= : \partial_z^n a(z) b(z) : / n!. \end{aligned}$$

Here $:$ stands for the *normal ordered product* of fields defined by

$$(1.3) \quad : a(z) b(z) : = a(z)_+ b(z) + b(z) a(z)_-,$$

where

$$a(z)_+ = \sum_{j \leq -1} a_{(j)} z^{-j-1}, \quad a(z)_- = \sum_{j \geq 0} a_{(j)} z^{-j-1}.$$

Definition 1.1. Let $(V, |0\rangle)$ be a *pointed vector space*, i.e., a vector space V with a fixed non-zero vector $|0\rangle \in V$, which will be referred to as the *vacuum vector*. A *state-field correspondence* is a linear map

$$Y: V \rightarrow glf(V), \quad a \mapsto Y(a, z) = \sum_{n \in \mathbb{Z}} a_{(n)} z^{-n-1},$$

such that following axioms hold:

$$(vacuum \ axioms) \quad Y(|0\rangle, z) = I_V, \quad Y(a, z)|0\rangle = a + T(a)z + \cdots \in V[[z]],$$

where $I_V \in \text{End } V$ is the identity operator, $T \in \text{End } V$,

$$(translation \ invariance) \quad [T, Y(a, z)] = Y(Ta, z) = \partial_z Y(a, z).$$

The linear operator T on V is called the *translation operator*.

Example 1.2. Let V be a unital algebra with a unit element $|0\rangle$, and let T be a derivation of V . Then

$$Y(a, z)b = (e^{zT}a)b, \quad a, b \in V,$$

is a state-field correspondence. In fact, all state-field correspondences for which the fields $Y(a, z)$ are formal power series in z are obtained in this way. We will call such a state-field correspondence *trivial*.

Example 1.3. Given two state-field correspondences $(V_i, |0\rangle_i, Y_i)$, $i = 1, 2$, one defines their tensor product $(V = V_1 \otimes V_2, |0\rangle = |0\rangle_1 \otimes |0\rangle_2, Y)$, where $Y(a_1 \otimes a_2, z) = Y(a_1, z) \otimes Y(a_2, z)$. This is again a state-field correspondence, the translation operator being $T = T_1 \otimes I + I \otimes T_2$.

Example 1.4. A special case of Example 1.3 is the tensor product of a state-field correspondence $(V, |0\rangle, Y)$ and a unital algebra $(A, 1)$, which is viewed as a trivial state-field correspondence with $T = 0$. In particular, the space $\text{Mat}_N(V)$ of N by N matrices with entries in V has a structure of a state-field correspondence.

Example 1.5. A generalization of Example 1.4 is the *smash product* $V \sharp \Gamma$, where Γ is a group of automorphisms of the state-field correspondence $(V, |0\rangle, Y)$. We define the space of states to be $V \otimes \mathbb{C}[\Gamma]$, the vacuum vector to be $|0\rangle \otimes 1$, and let

$$Y(a \otimes g, z)(b \otimes h) = Y(a, z)(gb) \otimes gh, \quad a, b \in V, g, h \in \Gamma.$$

When Γ is finite, one has the following very useful formula (cf. [MS]):

$$(1.4) \quad V^\Gamma \simeq \{e_{(-1)}(v_{(-1)}e) \mid v \in V \sharp \Gamma\} \subset V \sharp \Gamma, \quad \text{where } e = \frac{1}{|\Gamma|} \sum_{g \in \Gamma} |0\rangle \otimes g.$$

Here $V^\Gamma = \{a \in V \mid ga = a \text{ for all } g \in \Gamma\}$ denotes the subalgebra of invariants of Γ in V . The isomorphism (1.4) is given by $a \mapsto \frac{1}{|\Gamma|} \sum_{g \in \Gamma} a \otimes g$ for $a \in V^\Gamma$.

One defines in the obvious way the notions of subalgebras, ideals and homomorphisms of state-field correspondences. For example, a *subalgebra* of a state-field correspondence $(V, |0\rangle, Y)$ is a subspace U of V containing $|0\rangle$ and such that $a_{(n)}b \in U$ for all $n \in \mathbb{Z}$ if $a, b \in U$. A *left ideal* is a T -invariant subspace I of V such that $a_{(n)}b \in I$ for all $n \in \mathbb{Z}$ if $a \in V, b \in I$. If I is a two-sided ideal of V , there is a natural state-field correspondence with a space of states V/I .

Proposition 1.6. *In terms of Fourier coefficients $a_{(n)}$ of Y , the definition of a state-field correspondence can be reformulated as follows:*

$$\begin{aligned} & \text{(local finiteness)} \quad a_{(N)}b = 0 \quad \text{for } a, b \in V, N \gg 0, \\ & \text{(weak vacuum axiom)} \quad a_{(-1)}|0\rangle = |0\rangle_{(-1)}a = a, \\ & \text{(translation invariance)} \quad [T, a_{(n)}] = (Ta)_{(n)} = -na_{(n-1)} \quad \text{for } n \in \mathbb{Z}, a \in V. \end{aligned}$$

Proof. We have to check that $|0\rangle_{(n)}a = 0$ for $n \neq -1$, $Ta = a_{(-2)}|0\rangle$, and $a_{(n)}|0\rangle = 0$ for $n \geq 0$, $a \in V$. First, $Ta = (Ta)_{(-1)}|0\rangle = a_{(-2)}|0\rangle$. Next, we have $T|0\rangle = T(|0\rangle_{(-1)}|0\rangle) = 2T|0\rangle$, hence $T|0\rangle = 0$. Then for every $n \neq 0$, $(T|0\rangle)_{(n)} = -n|0\rangle_{(n-1)}$ shows that $|0\rangle_{(n-1)} = 0$. Similarly, $T(a_{(n)}|0\rangle) = -na_{(n-1)}|0\rangle$ implies $a_{(n)}|0\rangle = 0$ for $n \geq 0$, because $a_{(N)}|0\rangle = 0$ for $N \gg 0$. $\square$

Proposition 1.7. *Any state-field correspondence Y has the following properties:*

- (a) $Y(a, z)|0\rangle = e^{zT}a$,
- (b) $e^{wT}Y(a, z)e^{-wT} = Y(e^{wT}a, z) = i_{z,w}Y(a, z+w)$,
- (c) $(Y(a, z)_{(n)}Y(b, z))|0\rangle = Y(a_{(n)}b, z)|0\rangle$.

Proof. See [K, Proposition 4.1]. $\square$

Proposition 1.8. *Given a state-field correspondence Y , define*

$$(1.5) \quad Y^{\text{op}}(a, z)b = e^{zT}Y(b, -z)a.$$

Then Y^{op} is also a state-field correspondence, and $(Y^{\text{op}})^{\text{op}} = Y$.

Proof. Straightforward. $\square$

Definition 1.9. The state-field correspondence Y^{op} defined by (1.5) is called the *opposite* to Y .

At the end of this section, we study the notion of grading for state-field correspondences.

Definition 1.10. A state-field correspondence Y on a pointed vector space $(V, |0\rangle)$ is called *graded* if there is a diagonalizable operator $H \in \text{End } V$ satisfying

$$(1.6) \quad [H, Y(a, z)] = Y(Ha, z) + z\partial_z Y(a, z), \quad a \in V.$$

It is called $\mathbb{Z}_+$ -*graded* if all eigenvalues of H are non-negative integers. The operator H is called a *Hamiltonian* of V .

For a homogeneous element $a \in V$, we denote its degree by Δ_a : $Ha = \Delta_a a$. In terms of modes, equation (1.6) is equivalent to:

$$(1.7) \quad [H, a_{(n)}] = (\Delta_a - 1 - n)a_{(n)}, \quad a \in V, \quad Ha = \Delta_a a, \quad n \in \mathbb{Z}.$$

This implies that for homogeneous $a, b \in V$ one has:

$$(1.8) \quad \Delta_{a_{(n)}b} = \Delta_a + \Delta_b - n - 1, \quad n \in \mathbb{Z}.$$

One can easily show that $H|0\rangle = 0$ and $[H, T] = T$. The latter is equivalent to: $\Delta_{Ta} = \Delta_a + 1$ for homogeneous $a \in V$ (and is a special case of (1.8) since $Ta = a_{-2}|0\rangle$).

Proposition 1.11. *If Y is a graded state-field correspondence, then its opposite Y^{op} is also graded, with the same Hamiltonian H .*

Proof. Easy exercise, using (1.5), (1.6) and $He^{zT} = e^{zT}(H + zT)$. $\square$

Remark 1.12. A field $\phi(z) \in \text{glf}(V)$ is said to have *conformal dimension* Δ if

$$[H, \phi(z)] = (\Delta + z\partial_z)\phi(z).$$

Then for each homogeneous $a \in V$, $Y(a, z)$ is a field of conformal dimension Δ_a (see (1.6)).

Let $\phi(z), \phi'(z) \in \text{glf}(V)$ be two fields of conformal dimensions Δ and Δ' , respectively. Then $\partial_z \phi(z)$ has conformal dimension $\Delta + 1$, and for each $n \in \mathbb{Z}$, the n -th product $\phi(z)_{(n)}\phi'(z)$ has conformal dimension $\Delta + \Delta' - n - 1$.

If Y is a $\mathbb{Z}_+$ -graded state-field correspondence, we may shift subscripts as follows:

$$(1.9) \quad a_n = a_{(n+\Delta_a-1)}, \quad a_{(n)} = a_{n-\Delta_a+1} \quad \text{if } Ha = \Delta_a a,$$

so that

$$(1.10) \quad [H, a_n] = -n a_n, \quad a \in V, \quad n \in \mathbb{Z}.$$

It is easy to see that:

$$(1.11) \quad ((T + H)a)_0 = 0 \quad \text{for all } a \in V.$$

2. LOCALITY

A pair of fields $a(z), b(z) \in glf(V)$ is called *local* if

$$(z - w)^N [a(z), b(w)] = 0 \quad \text{for } N \gg 0.$$

It is called *local on* $v \in V$ if

$$(z - w)^N [a(z), b(w)] v = 0 \quad \text{for } N \gg 0.$$

Proposition 2.1. *A pair of fields $(a(z), b(z))$ is local if and only if*

$$[a(z), b(w)] = \sum_{\substack{j \geq 0 \\ \text{finite}}} (a(w)_{(j)} b(w)) \partial_w^j \delta(z - w) / j!.$$

It is local on v *iff*

$$[a(z), b(w)] v = \sum_{\substack{j \geq 0 \\ \text{finite}}} (a(w)_{(j)} b(w)) v \partial_w^j \delta(z - w) / j!.$$

Proof. Follows from [K, Corollary 2.2]. □

A pair of fields $(a(z), b(z))$ is called *weakly local* if

$$\text{Res}_z (z - w)^N [a(z), b(w)] = 0 \quad \text{for } N \gg 0.$$

Note that the weak locality of the pair $(a(z), b(z))$ means that $a(z)_{(N)} b(z) = 0$ for $N \gg 0$.

Remark 2.2. If a pair of fields $(a(z), b(z))$ is local, then the pair $(b(z), a(z))$ is also local. However, the weak locality of a pair (a, b) does not imply the weak locality of the pair (b, a) . Indeed, let $a(z)$ be the free boson, so that $[a_{(m)}, a_{(n)}] = m \delta_{m, -n}$, and take $b(z) = a_{(1)} z^{-1}$. Then $[a(z), b(w)] = -w^{-1}$; therefore, $a(z)_{(n)} b(z) = 0$, $b(z)_{(n)} a(z) = (-z)^n$ for all $n \geq 0$.

A collection of fields is called *local* (respectively *local on* $v \in V$, respectively *weakly local*) if each pair of fields from this collection is local (respectively local on $v \in V$, respectively weakly local).

A state-field correspondence Y is called *local* (respectively *weakly local*) if the collection of fields $\{Y(a, z)\}_{a \in V}$ is local (respectively weakly local).

Example 2.3. Let U be a vector space and let $V \subset glf(U)$ be a weakly local space of $\text{End } U$ -valued fields in the variable x . Let $|0\rangle = I_U \in V$. Then the following formula defines a state-field correspondence:

$$Y(a(x), z) b(x) = \sum_{n \in \mathbb{Z}} a(x)_{(n)} b(x) z^{-n-1}, \quad a(x), b(x) \in V,$$

the translation operator being $T = \partial_x$.

The following proposition generalizes the Uniqueness theorem from [K, Sec. 4.4].

Proposition 2.4. *Let Y be a state-field correspondence on a pointed vector space $(V, |0\rangle)$. Let $B_i(z) \in glf(V)$, $i = 1, 2$, be such that*

- (i) $B_1(z)|0\rangle = B_2(z)|0\rangle$,
- (ii) *all pairs $(Y(a, z), B_i(z))$ are local on $|0\rangle$.*

Then $B_1(z) = B_2(z)$.

Proof. Let $B(z) = B_1(z) - B_2(z)$. Then $B(z)|0\rangle = 0$ and all pairs $(Y(a, z), B(z))$ are local on $|0\rangle$. We have $(z-w)^N[B(z), Y(a, w)]|0\rangle = 0$, so $(z-w)^N B(z)Y(a, w)|0\rangle = 0$ for $N \gg 0$. Using Proposition 1.7a, we get $(z-w)^N B(z)e^{wT}a = 0$. Letting $w = 0$, we obtain $B(z)a = 0$, $a \in V$. $\square$

The following is a generalization of Dong's lemma (see [K, Lemma 3.2]).

Lemma 2.5. *Let $a, b, c \in \mathfrak{glf}(V)$ be three fields.*

- (a) *If the pair (a, b) is weakly local and the pairs (a, c) , (b, c) are local, then $(a_{(n)}b, c)$ is local for any $n \in \mathbb{Z}$.*
- (b) *If the pairs (a, b) and (a, c) are weakly local and the pair (b, c) is local, then $(a_{(n)}b, c)$ is local for any $n \geq 0$.*
- (c) *If the pairs (a, b) , (a, c) , (b, c) are weakly local, then $(a_{(n)}b, c)$ is weakly local for any $n \geq 0$.*
- (d) *If the pairs (a, b) , (a, c) , (b, c) are weakly local, then $(a, b_{(n)}c)$ is weakly local for any $n \in \mathbb{Z}$.*

Proof. Same as in [K, Sec. 3.2]. $\square$

Remark 2.6. In general, it is not true that if all pairs (a, b) , (a, c) , (b, c) are weakly local, then $(a_{(-1)}b, c)$ is weakly local as well. For example, let $a(z)$ be the free boson (see Remark 2.2), and take $b(z) = a(z)$, $c(z) = a(z)_+$. Then $[a(z)_-, a(w)_+] = i_{z,w}(z-w)^{-2}$ and $[a(z)_+, a(w)_+] = 0$; hence (a, a) and (a, a_+) are weakly local. We have: $[a(z)^2, a(w)_+] = [a(z)_+a(z) + a(z)a(z)_-, a(w)_+] = 2a(z)i_{z,w}(z-w)^{-2}$, which shows that the pair (a^2, a_+) is not weakly local.

Lemma 2.7. *Let X and Y be two state-field correspondences on a pointed vector space $(V, |0\rangle)$, and let $a, b, c \in V$ be such that*

- (i) *$(Y(a, z), Y(b, z))$ is a weakly local pair,*
- (ii) *the pair $(Y(a, z), X(c, z))$ (respectively, $(Y(b, z), X(c, z))$) is local on $|0\rangle$ and b (respectively, on $|0\rangle$ and a).*

Then the pair $(Y(a, z)_{(n)}Y(b, z), X(c, z))$ is local on the vacuum vector for all $n \in \mathbb{Z}$.

Proof. Let us write $a(z) = Y(a, z)$, $b(z) = Y(b, z)$, $c(z) = X(c, z)$ for short. By the (weak) locality, there exists a number $N \geq 0$ such that for all $r \geq N$:

$$\begin{aligned} \text{Res}_{z_1}(z_1 - z_2)^r[a(z_1), b(z_2)] &= 0, \\ (z_1 - z_3)^r[a(z_1), c(z_3)]v &= 0, \quad v = |0\rangle \text{ or } b, \\ (z_2 - z_3)^r[b(z_2), c(z_3)]v &= 0, \quad v = |0\rangle \text{ or } a. \end{aligned}$$

Let us apply the differential operator $e^{-z_2(\partial_{z_1} + \partial_{z_3})}$ to the left-hand side of the second equation for $v = b$. By Taylor's formula, the result is:

$$(z_1 - z_3)^r i_{z_1, z_2} i_{z_3, z_2} [a(z_1 - z_2), c(z_3 - z_2)]b.$$

Using Proposition 1.7, this is equal to:

$$\begin{aligned} &(z_1 - z_3)^r [e^{-z_2T}a(z_1)e^{z_2T}, e^{-z_2T}c(z_3)e^{z_2T}]b \\ &= (z_1 - z_3)^r e^{-z_2T} [a(z_1), c(z_3)] e^{z_2T} b \\ &= (z_1 - z_3)^r e^{-z_2T} [a(z_1), c(z_3)] b(z_2)|0\rangle. \end{aligned}$$

Applying $e^{z_2 T}$ to this, we obtain:

$$(z_1 - z_3)^r [a(z_1), c(z_3)] b(z_2) |0\rangle = 0, \quad r \geq N.$$

Similarly, we have:

$$(z_2 - z_3)^r [b(z_2), c(z_3)] a(z_1) |0\rangle = 0, \quad r \geq N.$$

The rest of the proof is as in [K, Lemma 3.2]. $\square$

In the proof of Lemma 2.7, we proved the following result which will be needed later.

Lemma 2.8. *Let X and Y be two state-field correspondences, and let $a, b, c \in V$ be such that*

$$(z - w)^N [Y(a, z), X(c, z)] b = 0$$

for some $N \geq 0$. Then

$$(z - w)^N [Y(a, z), X(c, z)] T b = 0.$$

Proposition 2.9. *Let X and Y be two state-field correspondences for $(V, |0\rangle)$, such that*

- (i) *Y is weakly local,*
- (ii) *all pairs $(Y(a, z), X(b, z))$ are local on any $v \in V$.*

Then $Y(a, z)_{(n)} Y(b, z) = Y(a_{(n)} b, z)$ for all $n \in \mathbb{Z}$, $a, b \in V$.

Proof. Let $B_1(z) = Y(a, z)_{(n)} Y(b, z)$, $B_2(z) = Y(a_{(n)} b, z)$. Due to Proposition 1.7c, the fields $B_i(z)$ satisfy condition (i) of Proposition 2.4. Due to Lemma 2.7, condition (ii) of Proposition 2.4 is satisfied as well. Hence $B_1(z) = B_2(z)$. $\square$

Proposition 2.10. *Let X and Y be two state-field correspondences. Then all pairs $(Y(a, z), X(b, z))$ are local on the vacuum vector if and only if $X = Y^{\text{op}}$.*

Proof. Assume that $(z - w)^N Y(a, z) X(b, w) |0\rangle = (z - w)^N X(b, w) Y(a, z) |0\rangle$ for $N \gg 0$. Using Proposition 1.7, this is equivalent to: $(z - w)^N e^{wT} i_{z,w} Y(a, z - w) b = (z - w)^N e^{zT} i_{w,z} X(b, w - z) a$. For a sufficiently large N , both sides contain only non-negative powers of $z - w$, and hence only non-negative powers of z and w . Putting $z = 0$, we obtain $X(b, w) a = e^{wT} Y(a, -w) b$.

Conversely, if $X = Y^{\text{op}}$, the above calculations show that $[Y(a, z), X(b, w)] |0\rangle = 0$ for all $a, b \in V$. $\square$

3. FIELD ALGEBRAS

Let $(V, |0\rangle)$ be a pointed vector space and let Y be a state-field correspondence. We say that Y satisfies the *n -th product axiom* if for all $a, b \in V$ and $n \in \mathbb{Z}$

$$(3.1) \quad Y(a_{(n)} b, z) = Y(a, z)_{(n)} Y(b, z).$$

We say that Y satisfies the *associativity axiom* if for all $a, b, c \in V$

$$(3.2) \quad (z - w)^N Y(Y(a, z) b, -w) c = (z - w)^N i_{z,w} Y(a, z - w) Y(b, -w) c, \quad N \gg 0.$$

Proposition 3.1. *Let $(V, |0\rangle)$ be a pointed vector space and let Y be a state-field correspondence. Then:*

(a) Y satisfies the n -th product axiom (3.1) iff for all $a, b \in V$

$$(3.3) \quad [Y(a, z), Y^{\text{op}}(b, w)] = \sum_{\substack{j \geq 0 \\ \text{finite}}} Y^{\text{op}}(a_{(j)}b, w) \partial_w^j \delta(z - w)/j!,$$

where $Y(a, z) = \sum_{j \in \mathbb{Z}} a_{(j)} z^{-j-1}$ and Y^{op} is the opposite to Y (see (1.5)).

(b) Y satisfies the associativity axiom (3.2) iff all pairs $(Y(a, z), Y^{\text{op}}(b, z))$ are local on each $v \in V$. In particular, the n -th product axiom implies the associativity axiom.

Proof. Assume that Y satisfies the n -th product axiom. Replace z with $-w$ in (3.1), multiply both sides by z^{-n-1} , and sum over $n \in \mathbb{Z}$. Then after some manipulation, we get:

$$(3.4) \quad \begin{aligned} & Y(Y(a, z)b, -w)c \\ &= i_{z,w} Y(a, z - w)Y(b, -w)c - Y(b, -w) \sum_{j \geq 0} (a_{(j)}c) \partial_w^j \delta(z - w)/j!. \end{aligned}$$

After applying e^{wT} to both sides, the left-hand side becomes $Y^{\text{op}}(c, w)Y(a, z)b$. The first term in the right-hand side becomes (using Proposition 1.7):

$$i_{z,w} e^{wT} Y(a, z - w)Y(b, -w)c = Y(a, z) e^{wT} Y(b, -w)c = Y(a, z) Y^{\text{op}}(c, w)b,$$

while the second term becomes:

$$\sum_{j \geq 0} Y^{\text{op}}(a_{(j)}c, w)b \partial_w^j \delta(z - w)/j!.$$

Therefore (3.4) is equivalent to (3.3) (with b replaced with c). Since (3.4) is the generating function of all n -th products, we get statement (a).

Notice that, by the above argument, the locality of the pair $(Y(a, z), Y^{\text{op}}(c, z))$ on b is equivalent to the associativity property (3.2). This completes the proof. $\square$

Corollary 3.2. *Let Y be a state-field correspondence for $(V, |0\rangle)$. Then for three elements $a, b, c \in V$, the collection of identities*

$$(3.5) \quad Y(a_{(n)}b, z)c = (Y(a, z)_{(n)}Y(b, z))c, \quad n \in \mathbb{Z}$$

implies the locality of the pair $(Y(a, z), Y^{\text{op}}(c, z))$ on b , which in turn is equivalent to the associativity property (3.2). If (3.5) holds for fixed a, c and all $b \in V$, then the pair $(Y(a, z), Y^{\text{op}}(c, z))$ is local.

Definition 3.3. Let $(V, |0\rangle)$ be a pointed vector space. A *field algebra* $(V, |0\rangle, Y)$ is a state-field correspondence Y for $(V, |0\rangle)$ satisfying the associativity axiom (3.2). A *strong field algebra* $(V, |0\rangle, Y)$ is a state-field correspondence Y satisfying the n -th product axiom (3.1).

Note that, by Proposition 3.1b, any strong field algebra is a field algebra.

Example 3.4. Let V be an associative algebra with a unit element $|0\rangle$ and let T be a derivation of V . Let $Y(a, z)b = (e^{zT}a)b$. Then $(V, |0\rangle, Y)$ is a strong field algebra. All field algebras for which all $Y(a, z)$ are formal power series in z are obtained in this way. We call such field algebras *trivial*.

Example 3.5. The tensor product of two field algebras (cf. Example 1.3) is again a field algebra. However, the tensor product of two strong field algebras is not necessarily a strong field algebra. For example, the tensor product of a non-trivial strong field algebra and a trivial field algebra (see Example 3.4), where the underlying associative algebra is non-commutative, is not a strong field algebra.

Example 3.6. The smash product of a field algebra V and a group Γ of its automorphisms (see Example 1.5) is a field algebra. It is not a strong field algebra if V is non-trivial and the action of Γ on V is non-trivial.

Theorem 3.7. (a) *A field algebra $(V, |0\rangle, Y)$ is the same as a state-field correspondence Y for $(V, |0\rangle)$ such that there exists a state-field correspondence X for $(V, |0\rangle)$, having the property that all pairs $(Y(a, z), X(b, z))$ are local on each $v \in V$. In this case $X = Y^{\text{op}}$.*
 (b) *A strong field algebra is the same as a field algebra $(V, |0\rangle, Y)$ for which the state-field correspondence Y is weakly local.*

Proof. (a) Let $(V, |0\rangle, Y)$ be a field algebra. Then, by Proposition 3.1b, Y^{op} is local with Y on each $v \in V$.

Conversely, let X be a state-field correspondence which is local with Y on every vector. In particular, it is local on the vacuum. Then, Proposition 2.10 implies that $X = Y^{\text{op}}$, and again by Proposition 3.1b, $(V, |0\rangle, Y)$ is a field algebra.

(b) If $(V, |0\rangle, Y)$ is a strong field algebra, then it is also a field algebra. The weak locality of Y follows from (3.1), because $a_{(n)}b = 0$ for $n \gg 0$.

Conversely, if $(V, |0\rangle, Y)$ is a field algebra with a weakly local Y , the n -th product axiom (3.1) follows from part (a) and Proposition 2.9. $\square$

Remark 3.8. If $(V, |0\rangle, Y)$ is a field algebra, then $(V, |0\rangle, Y^{\text{op}})$ is also a field algebra, called the *opposite* field algebra.

Corollary 3.9. *Let $(V, |0\rangle, Y)$ be a strong field algebra. Then for all $a, b \in V$ the pair $(Y(a, z), Y^{\text{op}}(b, z))$ is local and*

$$Y(a, z)_{(n)}Y^{\text{op}}(b, z) = Y^{\text{op}}(a_{(n)}b, z), \quad n \geq 0.$$

Proof. This follows from Proposition 3.1a. $\square$

Proposition 3.10. *A strong field algebra is a vector space V with a given vector $|0\rangle$ and a linear map $Y: V \rightarrow \text{glf}(V)$, $a \mapsto Y(a, z) = \sum_{n \in \mathbb{Z}} a_{(n)}z^{-n-1}$, satisfying for all $n \in \mathbb{Z}$, $a, b \in V$ the n -th product axiom (3.1) and the weak vacuum axioms*

$$(3.6) \quad |0\rangle_{(n)}a = \delta_{n,-1}a, \quad a_{(-1)}|0\rangle = a.$$

Proof. The only thing that has to be checked is that such a map Y is a state-field correspondence. Let $Ta = a_{(-2)}|0\rangle$. Letting $b = |0\rangle$ in n -th product axioms for $n \geq 1$, we see that the vacuum axioms hold. Notice that, taking coefficient of w^{-k-1} , the n -th product axiom gives the following relation:

$$(3.7) \quad (a_{(n)}b)_{(k)}c = \sum_{j=0}^{\infty} (-1)^j \binom{n}{j} (a_{(n-j)}(b_{(k+j)}c) - (-1)^n b_{(n+k-j)}(a_{(j)}c)).$$

Letting $k = -2$ and $c = |0\rangle$ in (3.7) gives $[T, a_{(n)}] = -n a_{(n-1)}$. Letting $n = -2$ and $b = |0\rangle$ in (3.7), we get $(Ta)_{(k)} = -k a_{(k-1)}$. Therefore, Y is a state-field correspondence. $\square$

Definition 3.11. A (strong) field algebra $(V, |0\rangle, Y)$ is called *graded*, respectively $\mathbb{Z}_+$ -*graded*, if the state-field correspondence Y is (see Definition 1.10).

Remark 3.12. Let $(V, |0\rangle, Y)$ be a $\mathbb{Z}_+$ -graded strong field algebra, and let $a, b, c \in V$ be homogeneous elements. Then $Y(a, z)_{(n)}Y(b, z) = 0$ for $n \geq \Delta_a + \Delta_b$, and the associativity relation (3.2) holds for $N \geq \Delta_a + \Delta_c$. This follows from the proof of Proposition 3.1 and (1.8).

Question 3.13. Is it true that a weakly local subspace V of $g\ell f(U)$, containing I_U , ∂_x -invariant and closed under all n -th products, is a strong field algebra? (See Example 2.3.)

4. CONFORMAL ALGEBRAS AND FIELD ALGEBRAS

Let $(V, |0\rangle)$ be a pointed vector space and let Y be a state-field correspondence. For $a, b \in V$, we define their λ -product by the formula

$$a_\lambda b = \text{Res}_z e^{\lambda z} Y(a, z)b = \sum_{\substack{n \geq 0 \\ \text{finite}}} \lambda^n a_{(n)} b / n!.$$

We also have the (-1) -st product on V , which we denote as

$$a.b = \text{Res}_z z^{-1} Y(a, z)b = a_{(-1)}b.$$

The vacuum axioms for Y imply:

$$(4.1) \quad |0\rangle.a = a = a.|0\rangle,$$

while the translation invariance axiom shows that:

$$(4.2) \quad T(a.b) = (Ta).b + a.(Tb)$$

and

$$(4.3) \quad T(a_\lambda b) = (Ta)_\lambda b + a_\lambda(Tb), \quad (Ta)_\lambda b = -\lambda a_\lambda b$$

for all $a, b \in V$.

Conversely, if we are given a linear operator T , a λ -product and a $.$ -product on V , satisfying the above properties (4.1)–(4.3), we can reconstruct the state-field correspondence Y by the formulas:

$$(4.4) \quad Y(a, z)_+ b = (e^{zT} a).b, \quad Y(a, z)_- b = (a_{-\partial_z} b)(z^{-1}),$$

where $Y(a, z) = Y(a, z)_+ + Y(a, z)_-$. Notice that equations (4.1)–(4.3) imply $T|0\rangle = 0$ and $|0\rangle_\lambda a = 0 = a_\lambda|0\rangle$ for $a \in V$ (cf. Proposition 1.6).

A $\mathbb{C}[T]$ -module V , equipped with a linear map $V \otimes V \rightarrow \mathbb{C}[\lambda] \otimes V$, $a \otimes b \mapsto a_\lambda b$, satisfying (4.3) is called a $(\mathbb{C}[T])$ -conformal algebra (cf. [K, Sec. 2.7]). On the other hand, with respect to the $.$ -product, V is a $(\mathbb{C}[T])$ -differential algebra (i.e., an algebra with a derivation T) with a unit $|0\rangle$.

We summarize the above discussion in the following lemma.

Lemma 4.1. *Giving a state-field correspondence on a pointed vector space $(V, |0\rangle)$ is equivalent to providing V with a structure of a $\mathbb{C}[T]$ -conformal algebra and a structure of a $\mathbb{C}[T]$ -differential algebra with a unit $|0\rangle$.*

Next, we translate the n -th product axioms in terms of the λ - and $.$ -products.

Lemma 4.2. *Let $(V, |0\rangle)$ be a pointed vector space and let Y be a state-field correspondence. Fix $a, b, c \in V$. Then the collection of n -th product identities (3.5) for $n \geq 0$ implies*

$$(4.5) \quad (a_\lambda b)_{\lambda+\mu} c = a_\lambda (b_\mu c) - b_\mu (a_\lambda c),$$

$$(4.6) \quad a_\lambda (b.c) = (a_\lambda b).c + b.(a_\lambda c) + \int_0^\lambda (a_\lambda b)_\mu c \, d\mu.$$

The (-1) -st product identity $Y(a_{(-1)} b, z) c = (Y(a, z)_{(-1)} Y(b, z)) c$ implies

$$(4.7) \quad (a.b)_\lambda c = (e^{T\partial_\lambda} a).(b_\lambda c) + (e^{T\partial_\lambda} b).(a_\lambda c) + \int_0^\lambda b_\mu (a_{\lambda-\mu} c) \, d\mu,$$

$$(4.8) \quad (a.b).c - a.(b.c) = \left(\int_0^T d\lambda a \right).(b_\lambda c) + \left(\int_0^T d\lambda b \right).(a_\lambda c).$$

Proof. The collection of n -th product identities (3.5) for $n \geq 0$ is equivalent to:

$$(4.9) \quad Y(a_\lambda b, w) c = \text{Res}_z e^{\lambda(z-w)} [Y(a, z), Y(b, w)] c = e^{-\lambda w} [a_\lambda, Y(b, w)] c.$$

Taking $\text{Res}_w e^{(\lambda+\mu)w}$, we obtain (4.5). Taking $\text{Res}_w w^{-1}$, and using that $e^{-\lambda w} w^{-1} = w^{-1} + \int_0^{-\lambda} e^{\mu w} d\mu$, we get:

$$(a_\lambda b).c = a_\lambda (b.c) - b.(a_\lambda c) + \int_0^{-\lambda} [a_\lambda, b_\mu] c \, d\mu.$$

This, together with (4.5), implies (4.6) (after the substitution $\mu' = \lambda + \mu$ in the integral).

Due to (1.3) and (4.4), the (-1) -st product identity is equivalent to:

$$Y(a.b, z) c = (e^{zT} a).(Y(b, z) c) + Y(b, z)(a_{-\partial_z} c)(z^{-1}).$$

Taking $\text{Res}_z e^{\lambda z}$ and using integration by parts, we get:

$$\begin{aligned} (a.b)_\lambda c &= \text{Res}_z (e^{T\partial_\lambda} e^{\lambda z} a).(Y(b, z) c) + \text{Res}_z Y(b, z)(a_{\lambda-\partial_z} c)(e^{\lambda z} z^{-1}) \\ &= (e^{T\partial_\lambda} a).(b_\lambda c) + \text{Res}_z Y(b, z)(a_{\lambda-\partial_z} c) \left(z^{-1} + \int_0^\lambda e^{\mu z} d\mu \right) \\ &= (e^{T\partial_\lambda} a).(b_\lambda c) + \text{Res}_z (e^{\partial_z \partial_\lambda} Y(b, z))(a_\lambda c) z^{-1} + \int_0^\lambda \text{Res}_z Y(b, z)(a_{\lambda-\mu} c) e^{\mu z} d\mu, \end{aligned}$$

which implies (4.7), using translation invariance.

On the other hand, taking $\text{Res}_z z^{-1}$ of the (-1) -st product identity, we get:

$$\begin{aligned} (a.b).c &= \text{Res}_z z^{-1} (Y(a, z)_+ Y(b, z)_+ c + Y(a, z)_+ Y(b, z)_- c + Y(b, z)_+ Y(a, z)_- c) \\ &= a.(b.c) + \text{Res}_z z^{-1} ((e^{zT} - 1)a).(Y(b, z)_- c) + \text{Res}_z z^{-1} ((e^{zT} - 1)b).(Y(a, z)_- c) \\ &= a.(b.c) + \text{Res}_z \left(\int_0^T e^{\lambda z} d\lambda a \right).(Y(b, z)_- c) + \text{Res}_z \left(\int_0^T e^{\lambda z} d\lambda b \right).(Y(a, z)_- c), \end{aligned}$$

which proves (4.8). $\square$

Remark 4.3. Equation (4.9) is equivalent to the commutator formula

$$(4.10) \quad [a_{(m)}, b_{(n)}] c = \sum_{j=0}^{\infty} \binom{m}{j} (a_{(j)} b)_{(m+n-j)} c, \quad m \in \mathbb{Z}_+, n \in \mathbb{Z}.$$

Equation (4.5) is equivalent to the same formula for $m, n \in \mathbb{Z}_+$.

Identity (4.5) is called the (left) *Jacobi identity* (cf. [K, Sec. 2.7]). A conformal algebra satisfying this identity for all $a, b, c \in V$ is called a (left) *Leibniz conformal algebra*. Equation (4.6) is known as the “non-commutative” *Wick formula* (cf. [K, (3.3.12)]), while (4.8) is called the *quasi-associativity formula* (cf. [K, (4.8.5)]).

Notice that the right-hand side of (4.8) is symmetric with respect to a and b , hence (4.8) implies

$$(4.11) \quad a.(b.c) - b.(a.c) = (a.b - b.a).c.$$

An algebra satisfying (4.11) for all $a, b, c \in V$ is called *left-symmetric*. For such an algebra $a.b - b.a$ is a Lie algebra bracket.

Theorem 4.4. *Giving a strong field algebra structure on a pointed vector space $(V, |0\rangle)$ is the same as providing V with a structure of a Leibniz $\mathbb{C}[T]$ -conformal algebra and a structure of a $\mathbb{C}[T]$ -differential algebra with a unit $|0\rangle$, satisfying (4.6)–(4.8).*

Proof. If $(V, |0\rangle, Y)$ is a strong field algebra, then by the above discussion we can define a λ -product and a $\cdot$ -product on V satisfying all the requirements.

Conversely, given a λ -product and a $\cdot$ -product, we define a state-field correspondence Y by (4.4). In the proof of Lemma 4.2, we have seen that equations (4.5)–(4.8) are equivalent to the identities

$$\text{Res}_z(Y(a_{(n)}b, z) - Y(a, z)_{(n)}Y(b, z))F = 0, \quad a, b \in V, \ n \geq -1, \ F = e^{\lambda z} \text{ or } z^{-1}.$$

Using the translation invariance of Y and integration by parts, we see that this identity holds also with F replaced with $\partial_z F$. Hence it holds for all $F = z^k$, $k < 0$. For $F = e^{\lambda z}$, taking coefficients at powers of λ shows that it is satisfied also for $F = z^k$, $k \geq 0$. This implies the n -th product axioms for $n \geq -1$.

Replacing a with Ta and using translation invariance shows that, for $n \neq 0$, the n -th product axiom implies the $(n-1)$ -st product axiom. Therefore, the n -th product axioms hold for all $n \in \mathbb{Z}$, and $(V, |0\rangle, Y)$ is a strong field algebra. $\square$

Corollary 4.5. *Let $(V, |0\rangle)$ be a pointed vector space, let Y be a state-field correspondence, and define the λ -product and $\cdot$ -product on V as above. Let U be a T -invariant subspace of V .*

- (a) *If (4.5) and (4.6) hold for all $a, b \in U$, $c \in V$, then $Y(a, z)_{(n)}Y(b, z) = Y(a_{(n)}b, z)$ for all $n \geq 0$, $a, b \in U$.*
- (b) *If (4.7) and (4.8) hold for all $a, b \in U$, $c \in V$, then $Y(a, z)_{(n)}Y(b, z) = Y(a_{(n)}b, z)$ for all $n < 0$, $a, b \in U$.*

5. TENSOR ALGEBRA OVER A LEIBNIZ CONFORMAL ALGEBRA

In this section, we are going to use the notation of the previous one. Let R be a Leibniz $\mathbb{C}[T]$ -conformal algebra (see Section 4), and let $V = \mathcal{T}(R) = \bigoplus_{m=0}^{\infty} R^{\otimes m}$ be the tensor algebra over R (viewed as a vector space over $\mathbb{C}$). We extend the action of T on R to V so that it is a derivation of the tensor product. Let $|0\rangle$ be the element $1 \in \mathbb{C} \equiv R^{\otimes 0} \subset V$.

Theorem 5.1. *There exists a unique structure of a field algebra on $V = T(R)$ such that the restriction of the λ -product to $R \times R$ coincides with the λ -product of R , the restriction of the $\cdot$ -product to $R \times V$ coincides with the tensor product, and the n -th product axioms (3.1) hold for $a \in R$, $b \in V$.*

For $a \in R$, $C \in V$, we set $a.C = a \otimes C$. Next, we define $a_\lambda C$ inductively using the Wick formula (4.6) and starting from $a_\lambda |0\rangle = 0$:

$$(5.1) \quad a_\lambda(c \otimes C) = (a_\lambda c) \otimes C + c \otimes (a_\lambda C) + \int_0^\lambda (a_\lambda c)_\mu C \, d\mu, \quad a, c \in R, C \in V.$$

Lemma 5.2. *Formula (5.1) defines a representation of the Leibniz conformal algebra R on V , whose restriction to R coincides with the adjoint representation.*

Proof. For $a, b \in R$, $C \in V$, we have to check the translation invariance:

$$(Ta)_\lambda C = -\lambda a_\lambda C, \quad a_\lambda(TC) = (\lambda + T)(a_\lambda C),$$

and the Jacobi identity (cf. (4.5)):

$$(a_\lambda b)_{\lambda+\mu} C = a_\lambda(b_\mu C) - b_\mu(a_\lambda C).$$

Note that these identities hold when $C \in R$ by definition. Assuming that they hold for a fixed $C \in V$ and all $a, b \in R$, we are going to prove them for $a, b, c \otimes C$ for any $a, b, c \in R$.

The translation invariance is immediate. The Jacobi identity reduces to checking that

$$\begin{aligned} & \int_0^{\lambda+\mu} ((a_\lambda b)_{\lambda+\mu} c)_\nu C \, d\nu \\ &= \int_0^\lambda (a_\lambda(b_\mu c))_\nu C \, d\nu + \int_0^\lambda (a_\lambda c)_\nu (b_\mu C) \, d\nu + \int_0^\mu a_\lambda((b_\mu c)_\nu C) \, d\nu \\ & \quad - \int_0^\mu (b_\mu(a_\lambda c))_\nu C \, d\nu - \int_0^\mu (b_\mu c)_\nu (a_\lambda C) \, d\nu - \int_0^\lambda b_\mu((a_\lambda c)_\nu C) \, d\nu. \end{aligned}$$

By Jacobi identity, the third and fifth terms in the right-hand side combine to $\int_0^\mu (a_\lambda(b_\mu c))_{\lambda+\nu} C \, d\nu = \int_\lambda^{\lambda+\mu} (a_\lambda(b_\mu c))_\nu C \, d\nu$, which together with the first term gives $\int_0^{\lambda+\mu} (a_\lambda(b_\mu c))_\nu C \, d\nu$. Similarly, the sum of the other three terms is $-\int_0^{\lambda+\mu} (b_\mu(a_\lambda c))_\nu C \, d\nu$. Now the statement follows from the Jacobi identity for a, b, c . $\square$

We have defined a λ -product $a_\lambda C$ and a $\cdot$ -product $a.C = a \otimes C$ for $a \in R$, $C \in V$, satisfying the translation invariance, the Jacobi identity, and the Wick formula. By the results of Section 4 (see Lemma 4.1 and Corollary 4.5), this gives a linear map from R to $glf(V)$, $a \mapsto Y(a, z)$, such that $[T, Y(a, z)] = Y(Ta, z) = \partial_z Y(a, z)$ and $Y(a, z)_{(n)} Y(b, z) = Y(a_{(n)} b, z)$ for all $n \geq 0$, $a, b \in R$.

Next, we set $Y(|0\rangle, z) = I_V$, and define fields $Y(A, z) \in glf(V)$ inductively by the formula

$$(5.2) \quad Y(a \otimes A, z) = :Y(a, z)Y(A, z): \quad \text{for } a \in R, A \in V.$$

It is easy to check that Y is a state-field correspondence.

Lemma 5.3. *We have $Y(a_{(n)} B, z) = Y(a, z)_{(n)} Y(B, z)$ for all $a \in R$, $B \in V$, $n \in \mathbb{Z}$. In particular, all pairs $(Y(a, z), Y(B, z))$ are weakly local for $a \in R$, $B \in V$.*

Proof. By definition, $Y(a_{(-1)}B, z) = Y(a \otimes B, z) = :Y(a, z)Y(B, z):$, so the statement holds for $n = -1$ and hence for all $n < 0$. Thus, it suffices to show that

$$Y(a_\lambda B, z) = [Y(a, z)_\lambda Y(B, z)] = \text{Res}_x e^{\lambda(x-z)} [Y(a, x), Y(B, z)], \quad a \in R, B \in V.$$

We have already proved this when $B \in R$, while for $B = |0\rangle$ both sides are trivial.

Assuming the above identity is true for $B \in V$ and all $a \in R$, consider it for $b \otimes B$ where $b \in R$. Using this assumption and (5.1), we find:

$$\begin{aligned} Y(a_\lambda(b \otimes B), z) &= [Y(a, z)_\lambda Y(b, z)]_{(-1)} Y(B, z) + Y(b, z)_{(-1)} [Y(a, z)_\lambda Y(B, z)] \\ &\quad + \int_0^\lambda [[Y(a, z)_\lambda Y(b, z)]_\mu Y(B, z)] d\mu. \end{aligned}$$

But the right-hand side is equal to $[Y(a, z)_\lambda (Y(b, z)_{(-1)} Y(B, z))]$ by the Wick formula for arbitrary fields [K, (3.3.12)]. This completes the proof. $\square$

Lemma 5.4. *All pairs $(Y(A, z), Y^{\text{op}}(B, z))$ are local for $A, B \in V$.*

Proof. It follows from Lemma 5.3 and Corollary 3.2 that all pairs $(Y(a, z), Y^{\text{op}}(B, z))$ are local for $a \in R, B \in V$. By induction, assume that $(Y(A, z), Y^{\text{op}}(B, z))$ is local, and consider the pair $(Y(a \otimes A, z), Y^{\text{op}}(B, z))$ for $a \in R$. Recall that by definition $Y(a \otimes A, z) = Y(a, z)_{(-1)} Y(A, z)$. Since, by Lemma 5.3, the pair $(Y(a, z), Y(A, z))$ is weakly local, we can apply Lemma 2.5a to conclude that the pair $(Y(a, z)_{(-1)} Y(A, z), Y^{\text{op}}(B, z))$ is local. $\square$

Now Theorem 3.7a and Lemma 5.4 imply that the so defined $(V, |0\rangle, Y)$ is a field algebra. Uniqueness is clear by construction. This completes the proof of Theorem 5.1.

The field algebra $V = \mathcal{T}(R)$ has the following universality property. Let W be a strong field algebra and let $f: R \rightarrow W$ be a homomorphism of conformal algebras (i.e., $f(a_{(n)}b) = f(a)_{(n)}f(b)$ and $f(Ta) = Tf(a)$ for all $a, b \in R, n \in \mathbb{Z}_+$). Then there is a unique homomorphism of field algebras $\tilde{f}: V \rightarrow W$ such that $f = \tilde{f} \circ i$, where i is the embedding of R in V .

Remark 5.5. Although all pairs $(Y(a, z), Y(B, z))$ are weakly local for $a \in R, B \in V$ (see Lemma 5.3), in general $(V, |0\rangle, Y)$ is not a strong field algebra. This follows from the results of Section 7.

Remark 5.6. A conformal algebra R is called *graded* if there is a diagonalizable operator $H \in \text{End } R$ satisfying $[H, T] = T$ and (1.7) for $n \in \mathbb{Z}_+$ and homogeneous $a \in R$. Then the tensor algebra $\mathcal{T}(R)$ over a Leibniz conformal algebra R is graded if R is graded. Moreover, $\mathcal{T}(R)$ is $\mathbb{Z}_+$ -graded if R is $\mathbb{Z}_+$ -graded.

6. FIELD ALGEBRAS AND VERTEX ALGEBRAS

Recall that a *vertex algebra* $(V, |0\rangle, Y)$ is a pointed vector space $(V, |0\rangle)$ together with a local state-field correspondence Y (see e.g. [K]). In particular, any vertex algebra is a strong field algebra with $Y = Y^{\text{op}}$ (see Theorem 3.7).

Example 6.1. The state-field correspondence given by Example 1.2 is a vertex algebra iff the algebra V is commutative and associative.

Example 6.2. Let V be a subspace of $g\ell f(U)$ which contains I_U , is ∂_x -invariant and is closed under all n -th products. Then the state-field correspondence defined in Example 2.3 gives V a structure of a vertex algebra iff V is a local collection. This follows from [K, Proposition 3.2].

Theorem 6.3. *A vertex algebra is the same as a field algebra $(V, |0\rangle, Y)$ for which $Y = Y^{\text{op}}$.*

Proof. Let $(V, |0\rangle, Y)$ be a field algebra with $Y = Y^{\text{op}}$. Recall that (see Proposition 3.1) the associativity relation (3.2) is equivalent to:

$$(z - w)^N [Y(a, z), Y^{\text{op}}(c, w)] b = 0.$$

By Lemma 2.8, this relation also holds after replacing b with Tb . Replacing b with $e^{uT}b$ in (3.2) and using Proposition 1.7b, we get:

$$(6.1) \quad \begin{aligned} & (z - w)^N i_{z,u} i_{w,u} Y(Y(a, z - u)b, u - w)c \\ &= (z - w)^N i_{z,w} i_{w,u} Y(a, z - w)Y(b, u - w)c. \end{aligned}$$

There exists $P \in \mathbb{Z}_+$ such that $b_{(k)}c = 0$ for $k \geq P$. Hence the right-hand side of (6.1) multiplied by $(u - w)^P$ contains only non-negative powers of $u - w$, hence only non-negative powers of w . Therefore, multiplying (6.1) by $(u - w)^P$ and putting $w = 0$ gives:

$$(6.2) \quad \begin{aligned} & Y(a, z)Y(b, u)c \\ &= z^{-N} u^{-P} (z - w)^N (u - w)^P i_{z,u} i_{w,u} Y(Y(a, z - u)b, u - w)c|_{w=0}. \end{aligned}$$

Using again Proposition 1.7b and $Y = Y^{\text{op}}$, we compute:

$$i_{z,u} Y(a, z - u)b = Y(e^{-uT}a, z)b = e^{zT}Y(b, -z)e^{-uT}a = i_{z,u} e^{(z-u)T}Y(b, u - z)a.$$

Therefore,

$$(6.3) \quad i_{z,u} i_{w,u} Y(Y(a, z - u)b, u - w) = i_{z,u} i_{w,z} Y(Y(b, u - z)a, z - w).$$

We may take $N = P \gg 0$ in (6.2). Comparing (6.2) and (6.3), we see that $Y(b, u)Y(a, z)c$ is given by the right-hand side of (6.2) where $i_{z,u}$ is replaced with $i_{u,z}$. Hence, if $K \in \mathbb{Z}_+$ is such that $a_{(k)}b = 0$ for $k \geq K$, we have

$$(z - u)^K Y(a, z)Y(b, u)c = (z - u)^K Y(b, u)Y(a, z)c,$$

which is locality of $Y(a, z)$ and $Y(b, z)$. This proves that $(V, |0\rangle, Y)$ is a vertex algebra. $\square$

Corollary 6.4. *A state-field correspondence is local iff it is local on every vector.*

Remark 6.5. It follows from (6.2), (6.3) that for $N \gg 0$ and any $n \in \mathbb{Z}$:

$$(6.4) \quad \begin{aligned} & Y(a, z)Y(b, u) i_{z,u}(z - u)^n - Y(b, u)Y(a, z) i_{u,z}(z - u)^n \\ &= z^{-N} u^{-N} (z - w)^N (u - w)^N i_{w,u}(i_{z,u} - i_{u,z})(z - u)^n \\ &\quad \times Y(Y(a, z - u)b, u - w)c|_{w=0}. \end{aligned}$$

Using (1.2), we find

$$(i_{z,u} - i_{u,z})(z - u)^n Y(a, z - u)b = \sum_{m=0}^{\infty} (a_{(m+n)}b) \partial_u^m \delta(z - u)/m!.$$

Note that this sum is finite, because $a_{(m)}b = 0$ for $m \gg 0$. Therefore, for large enough N , we can put $w = 0$ in (6.4), and obtain:

$$(6.5) \quad \begin{aligned} & Y(a, z)Y(b, u) i_{z,u}(z-u)^n - Y(b, u)Y(a, z) i_{u,z}(z-u)^n \\ &= \sum_{m=0}^{\infty} Y(a_{(m+n)}b, u) \partial_u^m \delta(z-u)/m!. \end{aligned}$$

The collection of these identities for $n \in \mathbb{Z}$ is equivalent to the Borcherds identity [K, (4.8.1)]. In particular, taking Res_z of both sides of (6.5), we obtain the n -th product identity (3.1).

Remark 6.6. Note that formula (6.1) holds for any field algebra $(V, |0\rangle, Y)$. Let $K \in \mathbb{Z}_+$ be such that $a_{(k)}b = 0$ for $k \geq K$. After multiplication by $(z-u)^K$, the left-hand side of (6.1) contains only non-negative powers of $z-u$, hence only non-negative powers of z . Putting $z = 0$ and replacing u with $-u$, we obtain:

$$\begin{aligned} & i_{w,u}Y(Y(a, u)b, -u-w)c \\ &= (-w)^{-N} u^{-K} (z-w)^N (z+u)^K i_{z,w} i_{w,u} Y(a, z-w) Y(b, -u-w) c \Big|_{z=0}. \end{aligned}$$

Since in both sides $-u-w$ is expanded in non-negative powers of u , we can replace w with $-u-w$ and use Taylor's formula to get:

$$(6.6) \quad \begin{aligned} & Y(Y(a, u)b, w)c \\ &= i_{z,u+w} i_{w,u} (u+w)^{-N} u^{-K} (z+u+w)^N (z+u)^K \\ & \quad \times Y(a, z+(u+w)) Y(b, w) c \Big|_{z=0}. \end{aligned}$$

Remark 6.7. Let V be a field algebra. Given $a, b \in V$, denote by ab the $\mathbb{C}$ -span of all elements $a_{(n)}b$ ($n \in \mathbb{Z}$). More generally, if A and B are two subspaces of V , we define AB as the $\mathbb{C}$ -span of all elements $a_{(n)}b$ ($n \in \mathbb{Z}$, $a \in A$, $b \in B$). Formulas (6.2) and (6.6) imply associativity of this product:

$$(6.7) \quad (ab)c = a(bc), \quad a, b, c \in V.$$

This property allows one to apply many arguments of the theory of associative algebras to field algebras. Notice that (6.7) holds also for $a, b \in V$ and $c \in M$, where M is a module over the field algebra V (see Definition 8.1 below).

Remark 6.8. Letting $a \bullet b = \mathbb{C}[T](ab)$, we still have associativity of the product $\bullet$ for an arbitrary field algebra V . In addition, $|0\rangle \bullet a = a \bullet |0\rangle = \mathbb{C}[T]a$ for all $a \in V$. If V is a vertex algebra, the product $\bullet$ is also commutative. These remarks allow one to use arguments of commutative algebra to study vertex algebras (this will be pursued in a future publication).

For the opposite state-field correspondence Y^{op} , we can consider

$$a_{\lambda}^{\text{op}} b = \text{Res}_z e^{\lambda z} Y^{\text{op}}(a, z)b, \quad a^{\text{op}} b = \text{Res}_z z^{-1} Y^{\text{op}}(a, z)b,$$

and express these operations in terms of the λ -product and $\cdot$ -product defined for Y . We have:

$$(6.8) \quad a_{\lambda}^{\text{op}} b = \text{Res}_z e^{\lambda z} e^{zT} Y(b, -z)a = -\text{Res}_z e^{-(\lambda+T)z} Y(b, z)a = -b_{-\lambda-T} a,$$

and, using $z^{-1}e^{zT} = z^{-1} - \int_0^{-T} e^{-\lambda z} d\lambda$,

$$(6.9) \quad a^{\text{op}}b = \text{Res}_z z^{-1}e^{zT}Y(b, -z)a = b.a + \int_0^{-T} b_\lambda a d\lambda.$$

If $(V, |0\rangle, Y)$ is a vertex algebra, then the λ -product satisfies the *skewsymmetry relation* $a_\lambda b = -b_{-\lambda-T}a$ for $a, b \in V$. A conformal algebra satisfying the Jacobi identity and the skewsymmetry relation is called a *Lie conformal algebra* [K, Sec. 2.7].

Theorem 6.9. *Giving a vertex algebra structure on a pointed vector space $(V, |0\rangle)$ is the same as providing V with the structures of a Lie $\mathbb{C}[T]$ -conformal algebra and a left-symmetric $\mathbb{C}[T]$ -differential algebra with a unit $|0\rangle$ (see (4.11)), satisfying the Wick formula (4.6) and*

$$(6.10) \quad a.b - b.a = \int_{-T}^0 a_\lambda b d\lambda, \quad a, b \in V.$$

Proof. By Lemma 4.1, these data define a state-field correspondence Y . By (6.8) and (6.9), we have $Y = Y^{\text{op}}$; therefore, we only need to prove that V is a field algebra (see Theorem 6.3). Due to Theorem 4.4, it remains to show that (4.7) and (4.8) follow from equations (4.5), (4.6), (4.11), (6.10) and the skewsymmetry of the λ -product.

First, using (6.10), rewrite the Wick formula (4.6) as

$$a_\lambda(b.c) = c.(a_\lambda b) + b.(a_\lambda c) + \int_{-T}^\lambda (a_\lambda b)_\mu c d\mu.$$

Then replace λ with $-\lambda$ in this equation and apply $e^{T\partial_\lambda}$ to both sides. Using Taylor's formula and the fact that T is a derivation, we obtain:

$$a_{-\lambda-T}(b.c) = (e^{T\partial_\lambda} c).(a_{-\lambda-T} b) + (e^{T\partial_\lambda} b).(a_{-\lambda-T} c) + \int_{-T}^{-\lambda-T} e^{T\partial_\lambda} ((a_{-\lambda} b)_\mu c) d\mu.$$

Using (4.3) and the skewsymmetry of the λ -product, the last term can be rewritten as follows:

$$\begin{aligned} \int_{-T}^{-\lambda-T} e^{T\partial_\lambda} ((a_{-\lambda} b)_\mu c) d\mu &= - \int_{-T}^{-\lambda-T} e^{T\partial_\lambda} (c_{-\mu-T}(a_{-\lambda} b)) d\mu \\ &= \int_0^\lambda e^{T\partial_\lambda} (c_\mu(a_{-\lambda} b)) d\mu = \int_0^\lambda c_\mu(a_{-\lambda+\mu-T} b) d\mu. \end{aligned}$$

Applying the skewsymmetry relation again, we obtain (4.7).

In order to prove (4.8), we manipulate its left-hand side using twice (6.10):

$$(a.b).c - a.(b.c) = c.(a.b) - a.(c.b) + \int_{-T}^0 d\lambda (a.b)_\lambda c - a. \left(\int_{-T}^0 d\lambda b_\lambda c \right).$$

Due to (4.11) and (6.10), the first two terms in the right-hand side of this equation give:

$$\begin{aligned} c.(a.b) - a.(c.b) &= -(a.c - c.a).b = -\left(\int_{-T}^0 d\lambda a_\lambda c\right).b \\ &= -b.\left(\int_{-T}^0 d\lambda a_\lambda c\right) + \int_{-T}^0 d\mu b_\mu \left(\int_{-T}^0 d\lambda a_\lambda c\right). \end{aligned}$$

The double integral is equal to:

$$\begin{aligned} (6.11) \quad \int_{-T}^0 d\mu b_\mu \left(\int_{-T}^0 d\lambda a_\lambda c\right) &= \int_{-T}^0 d\mu \int_{-T-\mu}^0 d\lambda b_\mu(a_\lambda c) \\ &= \int_{-T}^0 d\mu \int_{-T}^\mu d\lambda b_\mu(a_{\lambda-\mu}c) = \int_{-T}^0 d\lambda \int_\lambda^0 d\mu b_\mu(a_{\lambda-\mu}c). \end{aligned}$$

Therefore, applying (4.7), which we already proved, we obtain:

$$\begin{aligned} (a.b).c - a.(b.c) &= \int_{-T}^0 d\lambda (e^{T\partial_\lambda} a).(b_\lambda c) - a.\left(\int_{-T}^0 d\lambda b_\lambda c\right) \\ &\quad + \int_{-T}^0 d\lambda (e^{T\partial_\lambda} b).(a_\lambda c) - b.\left(\int_{-T}^0 d\lambda a_\lambda c\right). \end{aligned}$$

From here, it is easy to deduce (4.8).

This completes the proof of the theorem. $\square$

Proposition 6.10. *If R is a Leibniz $\mathbb{C}[T]$ -conformal algebra with a λ -product $a_\lambda b$, then the bracket*

$$(6.12) \quad [a, b] = \int_{-T}^0 a_\lambda b d\lambda, \quad a, b \in R$$

defines the structure of a Leibniz $\mathbb{C}[T]$ -differential algebra on R . If R is a Lie conformal algebra, then this is a Lie bracket.

Proof. It is clear that T is a derivation of the bracket (6.12), because it is a derivation of the λ -product $a_\lambda b$. From relation (6.11), we have:

$$[b, [a, c]] = \int_{-T}^0 d\lambda \int_\lambda^0 d\mu b_\mu(a_{\lambda-\mu}c) = \int_{-T}^0 d\lambda \int_\lambda^0 d\mu b_{\lambda-\mu}(a_\mu c),$$

and similarly,

$$[a, [b, c]] = \int_{-T}^0 d\lambda \int_\lambda^0 d\mu a_\mu(b_{\lambda-\mu}c).$$

On the other hand,

$$[[a, b], c] = \int_{-T}^0 d\lambda \left(\int_{-T}^0 d\mu a_\mu b\right)_\lambda c = \int_{-T}^0 d\lambda \int_\lambda^0 d\mu (a_\mu b)_\lambda c.$$

Therefore, the Jacobi identity for the bracket (6.12) follows from the Jacobi identity (4.5) for the λ -product. Finally, it is easy to see that $[a, b] = -[b, a]$ if $a_\lambda b = -b_{-\lambda-T}a$. $\square$

Remark 6.11. Let R be a Lie $\mathbb{C}[T]$ -conformal algebra with a λ -product $a_\lambda b = \sum_{m \geq 0} \lambda^m a_{(m)} b / m!$. To it one can associate a Lie algebra $\text{Lie } R$ as follows (see [K, Sec. 2.7]). As a vector space, $\text{Lie } R = R[t, t^{-1}] / (T + \partial_t)R[t, t^{-1}]$. The Lie bracket in $\text{Lie } R$ is given by the formula (cf. (4.10)):

$$[a_{[m]}, b_{[n]}] = \sum_{j=0}^{\infty} \binom{m}{j} (a_{(j)} b)_{[m+n-j]}, \quad a, b \in R, \quad m, n \in \mathbb{Z},$$

where $a_{[m]}$ is the image of at^m in $\text{Lie } R$. Note that T acts on $\text{Lie } R$ as a derivation: $T(a_{[m]}) = -m a_{[m-1]}$.

Let $(\text{Lie } R)_-$ (respectively, $(\text{Lie } R)_+$) be the $\mathbb{C}$ -span of all $a_{[m]}$ for $a \in R$, $m \geq 0$ (respectively, $m < 0$). These are subalgebras of $\text{Lie } R$. Let R_{Lie} be R considered as a Lie algebra with respect to the bracket (6.12). Then the map $a \mapsto a_{[-1]}$ is an isomorphism of $\mathbb{C}[T]$ -differential Lie algebras $R_{\text{Lie}} \xrightarrow{\sim} (\text{Lie } R)_+$.

Theorem 6.12. *Let R be a Lie $\mathbb{C}[T]$ -conformal algebra with a λ -product $a_\lambda b$. Let R_{Lie} be R considered as a Lie algebra with respect to the bracket (6.12), and let $V = U(R_{\text{Lie}})$ be its universal enveloping algebra. Then there exists a unique structure of a vertex algebra on V such that the restriction of the λ -product to $R_{\text{Lie}} \times R_{\text{Lie}}$ coincides with the λ -product of R and the restriction of the $\cdot$ -product to $R_{\text{Lie}} \times V$ coincides with the product in $U(R_{\text{Lie}})$.*

Proof. We have $R_{\text{Lie}} \simeq (\text{Lie } R)_+$ (see Remark 6.11). By construction,

$$V \simeq U((\text{Lie } R)_+) \simeq \text{Ind}_{(\text{Lie } R)_-}^{\text{Lie } R} \mathbb{C}$$

(where $(\text{Lie } R)_-$ acts trivially on $\mathbb{C}$) coincides with the universal vertex algebra associated to $\text{Lie } R$ (see [K, Sec. 4.7]). $\square$

The vertex algebra V from Theorem 6.12 has the following universality property. Let W be another vertex algebra and let $f: R \rightarrow W$ be a homomorphism of conformal algebras (i.e., $f(a_{(n)} b) = f(a)_{(n)} f(b)$ and $f(Ta) = Tf(a)$ for all $a, b \in R$, $n \in \mathbb{Z}_+$). Then there is a unique homomorphism of vertex algebras $\tilde{f}: V \rightarrow W$ such that $f = \tilde{f} \circ i$, where i is the embedding of R in V . For this reason, V is called the *universal vertex algebra* associated to R and is denoted by $\mathcal{U}(R)$.

Remark 6.13. $\mathcal{U}(R)$ is the quotient of the field algebra $\mathcal{T}(R)$, constructed in Section 5, by the two-sided ideal generated by $a \otimes b - b \otimes a - \int_{-T}^0 a_\lambda b d\lambda$ ($a, b \in R$). A different proof of Theorem 6.12 was given in [GMS].

7. STRONG FIELD ALGEBRAS AND VERTEX ALGEBRAS

Let $(V, |0\rangle, Y)$ be a strong field algebra. We are going to use the λ -product and $\cdot$ -product on V defined in Section 4.

Fix four elements $a, a', b, c \in V$. Using formulas (4.6), (4.7), it is straightforward to compute:

$$(a.a')_\lambda(b_\mu c) = (e^{T\partial_\lambda} a).(a'_\lambda(b_\mu c)) + (e^{T\partial_\lambda} a').(a_\lambda(b_\mu c)) + \int_0^\lambda d\nu a'_\nu(a_{\lambda-\nu}(b_\mu c)),$$

$$\begin{aligned} b_\mu((a.a')_\lambda c) &= (e^{T\partial_\lambda} a).(b_\mu(a'_\lambda c)) + (e^{T\partial_\lambda}(b_\mu a)).(a'_{\lambda+\mu} c) \\ &\quad + \int_0^\mu d\nu (b_\mu a)_\nu(a'_{\lambda+\mu-\nu} c) + (a \leftrightarrow a') + \int_0^\lambda d\nu b_\mu(a'_\nu(a_{\lambda-\nu} c)), \end{aligned}$$

and

$$\begin{aligned} ((a.a')_\lambda b)_{\lambda+\mu} c &= (e^{T\partial_\lambda} a).((a'_\lambda b)_{\lambda+\mu} c) + (e^{T\partial_\lambda}(a'_{-\mu-T} b)).(a_{\lambda+\mu} c) \\ &\quad + \int_0^{\lambda+\mu} d\nu (a'_{\nu-\mu} b)_\nu(a_{\lambda+\mu-\nu} c) + (a \leftrightarrow a') + \int_0^\lambda d\nu (a'_\nu(a_{\lambda-\nu} b))_{\lambda+\mu} c, \end{aligned}$$

where $(a \leftrightarrow a')$ denotes the sum of the first three terms with a and a' interchanged. Using Jacobi identity (4.5) for elements of the form $u, u_{(m)}v$ for $u, v \in \{a, a', b, c\}$, $m \in \mathbb{Z}_+$, after a long calculation one obtains:

$$\begin{aligned} (7.1) \quad &((a.a')_\lambda b)_{\lambda+\mu} c - (a.a')_\lambda(b_\mu c) + b_\mu((a.a')_\lambda c) \\ &= (e^{T\partial_\lambda}(a_{-\mu-T} b + b_\mu a)).(a'_{\lambda+\mu} c) + (a \leftrightarrow a'). \end{aligned}$$

This implies the following lemma.

Lemma 7.1. *Let V be a vector space equipped with a structure of a $\mathbb{C}[T]$ -conformal algebra and a structure of a $\mathbb{C}[T]$ -differential algebra. Let $a, a', b, c \in V$, and assume that formulas (4.5)–(4.7) hold for all elements of the form $u, u_{(m)}v$ ($u, v \in \{a, a', b, c\}$, $m \in \mathbb{Z}_+$). Then Jacobi identity (4.5) holds for the triple $(a.a', b, c)$ if and only if*

$$(7.2) \quad (a_{-\mu-T} b + b_\mu a).(a'_\lambda c) + (a \leftrightarrow a') = 0.$$

Proof. By the above calculation, Jacobi identity for $(a.a', b, c)$ is equivalent to the vanishing of the right-hand side of (7.1). Applying to it $e^{-T\partial_\lambda}$ and using Taylor's formula, we obtain:

$$(a_{-\mu-T} b + b_\mu a).(a'_{\lambda+\mu-T} c) + (a \leftrightarrow a') = 0,$$

which is equivalent to (7.2). $\square$

Applying first (4.7) and then (4.6), we find:

$$\begin{aligned} (a.a')_\lambda(b.c) &= (e^{T\partial_\lambda} a).((a'_\lambda b).c + b.(a'_\lambda c) + \int_0^\lambda d\mu (a'_\lambda b)_\mu c) + \int_0^\lambda d\mu (a_\mu b).(a'_{\lambda-\mu} c) \\ &\quad + (a \leftrightarrow a') + \int_0^\lambda d\mu \left\{ (a'_\mu(a_{\lambda-\mu} b)).c + b.(a'_\mu(a_{\lambda-\mu} c)) \right\} \\ &\quad + \int_0^\lambda d\mu \int_0^\mu d\nu \left\{ (a'_\mu(a_{\lambda-\mu} b))_\nu c + (a'_\mu b)_\nu(a_{\lambda-\mu} c) \right\} \\ &\quad + \int_0^\lambda d\mu \int_0^{\lambda-\mu} d\nu a'_\mu((a_{\lambda-\mu} b)_\nu c), \end{aligned}$$

where $(a \leftrightarrow a')$ denotes the sum of the first two terms with a and a' interchanged. We also have:

$$((a.a')_\lambda b).c = \left((e^{T\partial_\lambda} a).(a'_\lambda b) + (e^{T\partial_\lambda} a').(a_\lambda b) + \int_0^\lambda d\mu a'_\mu (a_{\lambda-\mu} b) \right).c$$

and

$$b.((a.a')_\lambda c) = b. \left((e^{T\partial_\lambda} a).(a'_\lambda c) + (e^{T\partial_\lambda} a').(a_\lambda c) + \int_0^\lambda d\mu a'_\mu (a_{\lambda-\mu} c) \right).$$

Using Jacobi identity (4.5) and quasi-associativity (4.8) for elements of the form $T^n u$, $u_{(m)} v$ for $u, v \in \{a, a', b, c\}$, $m, n \in \mathbb{Z}_+$, after a long calculation one can show that:

$$(7.3) \quad \begin{aligned} & (a.a')_\lambda (b.c) - ((a.a')_\lambda b).c - b.((a.a')_\lambda c) - \int_0^\lambda d\mu ((a.a')_\lambda b)_\mu c \\ &= \left((e^{T\partial_\lambda} a).b - b.(e^{T\partial_\lambda} a) - \int_{-T}^0 d\mu (e^{T\partial_\lambda} a)_\mu b \right).(a'_\lambda c) + (a \leftrightarrow a'). \end{aligned}$$

This implies:

Lemma 7.2. *Let V be a vector space equipped with a structure of a $\mathbb{C}[T]$ -conformal algebra and a structure of a $\mathbb{C}[T]$ -differential algebra. Let $a, a', b, c \in V$, and assume that formulas (4.5)–(4.8) hold for all elements of the form $T^n u$, $u_{(m)} v$ ($u, v \in \{a, a', b, c\}$, $m, n \in \mathbb{Z}_+$). Then the Wick formula (4.6) holds for $(a.a', b, c)$ if and only if*

$$(7.4) \quad \left(a.b - b.a - \int_{-T}^0 d\mu a_\mu b \right).(a'_\lambda c) + (a \leftrightarrow a') = 0.$$

Proof. By the above calculation, Wick formula for $(a.a', b, c)$ is equivalent to the vanishing of the right-hand side of (7.3). Note that (4.6) holds for (a, b, c) if and only if it holds for (Ta, b, c) . Replacing in (7.3) a and a' by $e^{-T\partial_\lambda} a$ and $e^{-T\partial_\lambda} a'$, respectively, we obtain (7.4). $\square$

The above two lemmas, together with Theorem 4.4 and relations (6.8), (6.9), immediately imply the following:

Corollary 7.3. *Let $(V, |0\rangle, Y)$ be a strong field algebra. Then all $a, a', b, c \in V$ satisfy*

$$(7.5) \quad \left(Y(a, z)b - Y^{\text{op}}(a, z)b \right).(a'_\lambda c) + (a \leftrightarrow a') = 0.$$

From this, we deduce the following surprising result.

Theorem 7.4. *Let $(V, |0\rangle, Y)$ be a strong field algebra such that $|0\rangle = a_{(n)}c$ for some $a, c \in V$, $n \in \mathbb{Z}_+$. Then V is a vertex algebra.*

In particular, if V contains a Virasoro element, i.e., an element ℓ such that $\ell_\lambda \ell = (T + 2\lambda)\ell + \alpha\lambda^3|0\rangle$, with $\alpha \in \mathbb{C} \setminus \{0\}$, then V is a vertex algebra.

Proof. Taking $a = a'$ and coefficient in front of λ^n in (7.5), we obtain $(Y(a, z)b - Y^{\text{op}}(a, z)b).|0\rangle = 0$, hence $Y(a, z) = Y^{\text{op}}(a, z)$. Then (7.5) implies $Y(a', z) = Y^{\text{op}}(a', z)$ for all $a' \in V$. By Theorem 6.3, V is a vertex algebra. $\square$

8. MODULES OVER FIELD ALGEBRAS

Definition 8.1. A *module* over a field algebra $(V, |0\rangle, Y)$ is a vector space M equipped with an operator $T^M \in \text{End } M$ and a linear map

$$V \rightarrow \text{glf}(M), \quad a \mapsto Y^M(a, z) = \sum_{n \in \mathbb{Z}} a_{(n)}^M z^{-n-1},$$

satisfying:

$$\begin{aligned} (\text{vacuum axiom}) \quad & Y^M(|0\rangle, z) = I_M, \\ (\text{translation invariance}) \quad & [T^M, Y^M(a, z)] = \partial_z Y^M(a, z), \\ (\text{associativity axiom}) \quad & (z-w)^N Y^M(Y(a, z)b, -w)v \\ & = (z-w)^N i_{z,w} Y^M(a, z-w) Y^M(b, -w)v \end{aligned}$$

for $a, b \in V$, $v \in M$, $N \gg 0$.

When $(V, |0\rangle, Y)$ is a strong field algebra, in the above definition we replace the associativity axiom with the following

$$(n\text{-th product axiom}) \quad Y^M(a_{(n)}b, z) = Y^M(a, z)_{(n)} Y^M(b, z), \quad n \in \mathbb{Z}.$$

Remark 8.2. What we introduced in Definition 8.1 are left modules over a field algebra. We can define a *right module* over a field algebra $(V, |0\rangle, Y)$ as a left module over the opposite field algebra $(V, |0\rangle, Y^{\text{op}})$. Any field algebra is both a left and a right module over itself.

Note that, as in Proposition 3.1, the n -th product axiom implies the associativity axiom. Taking $b = |0\rangle$ in the associativity axiom and using Proposition 1.7b, we deduce

$$(8.1) \quad Y^M(Ta, z) = \partial_z Y^M(a, z).$$

Remark 8.3. Let $(V, |0\rangle, Y)$ be a vertex algebra, and let M be a module over V considered as a field algebra. Then as in Section 6 one can show that (see Remark 6.5):

$$\begin{aligned} & Y^M(a, z) Y^M(b, u) i_{z,u}(z-u)^n - Y^M(b, u) Y^M(a, z) i_{u,z}(z-u)^n \\ (8.2) \quad & = \sum_{m=0}^{\infty} Y^M(a_{(m+n)}b, u) \partial_u^m \delta(z-u)/m!, \quad a, b \in V, n \in \mathbb{Z}. \end{aligned}$$

Therefore, M is also a module over V considered as a vertex algebra.

Let M be a module over a field algebra $(V, |0\rangle, Y)$. Using (8.1), the same argument as in Section 6 gives the following generalization of formula (6.2):

$$(8.3) \quad \begin{aligned} & Y^M(a, z) Y^M(b, u)v \\ & = z^{-N} u^{-P} (z-w)^N (u-w)^P i_{z,u} i_{w,u} Y^M(Y(a, z-u)b, u-w)v|_{w=0}, \end{aligned}$$

where $N \in \mathbb{Z}_+$ is from the associativity axiom and $P \in \mathbb{Z}_+$ is such that $b_{(k)}v = 0$ for $k \geq P$.

Formula (8.3) implies immediately the following lemma.

Lemma 8.4. *Let M be a module over a field algebra V and let $\mathcal{A}$ be the associative subalgebra of $\text{End}_{\mathbb{C}} M$ generated by all endomorphisms $a_{(n)}^M$ ($a \in V, n \in \mathbb{Z}$). Then for any $\alpha \in \mathcal{A}$, $v \in M$, αv is a $\mathbb{C}$ -linear combination of $a_{(n)}^M v$.*

This lemma along with Jacobson's density theorem (see e.g. [L]) implies:

Theorem 8.5 (Density theorem). *Let M be a semisimple module over a field algebra V (i.e., M is a direct sum of irreducible V -modules), let $K = \text{End}_V M$, $f \in \text{End}_K M$ and let U be a finite-dimensional (over $\mathbb{C}$) subspace of M . Then there exists a finite set of elements $\{a^i\}_{i \in I}$ of V and a set of integers $\{n_i\}_{i \in I}$ such that*

$$\sum_i a_{(n_i)}^{iM} |U = f|U.$$

In particular, if M is an irreducible countable-dimensional V -module, then the conclusion of the theorem holds for any $f \in \text{End}_{\mathbb{C}} M$.

Likewise, the proof of Theorem 1.1 from [KR] adapted to the field algebra framework (by making use of Theorem 8.5 in place of Jacobson's density theorem) gives the field algebra analogue of that theorem:

Theorem 8.6 (Duality theorem). *Let V be a field algebra and let M be an irreducible V -module. Let G be a group and suppose one has an action of G on V by automorphisms and a representation of G in M such that the following three properties hold:*

- (compatibility) $g(a_{(n)}v) = (ga)_{(n)}(gv)$, for $g \in G$, $a \in V$, $v \in M$, $n \in \mathbb{Z}$;
- (semisimplicity of G on V) V is a direct sum of irreducible G -modules;
- (semisimplicity of G on M) M is a direct sum of at most countable number of finite-dimensional irreducible G -modules.

Then each non-zero isotypic component M_π (where π is a finite-dimensional irreducible G -module) is irreducible with respect to the pair (V^G, G) (where V^G is the fixed point subalgebra of the field algebra V).

Remark 8.7. It was stated in [KT] that the proof of Theorem 1.1 from [KR] can be adapted to give Theorem 8.6 for vertex algebras. A proof in the case of graded vertex algebras with the use of Zhu's algebra was given in [DLM, DY].

Consider again formula (8.3), where $N \in \mathbb{Z}_+$ is from the associativity axiom and $P \in \mathbb{Z}_+$ is such that $b_{(k)}v = 0$ for $k \geq P$. Taking $\text{Res}_u u^{P-1}$, we obtain:

$$(8.4) \quad Y^M(a, z)(b_{(P-1)}^M v) = \sum_{i=0}^N \binom{N}{i} z^{-i} (-w)^{P+i} Y^M(Y(a, z)b, -w)v|_{w=0},$$

which implies

$$(8.5) \quad a_{(n)}^M(b_{(P-1)}^M v) = \sum_{i=0}^N \binom{N}{i} (a_{(n-i)}b)_{(P+i-1)}^M v, \quad n \in \mathbb{Z}.$$

Since the right-hand side of (8.4) contains only non-negative powers of w , we have:

$$(8.6) \quad 0 = \sum_{i=0}^N \binom{N}{i} (a_{(n-i)}b)_{(P+i)}^M v, \quad n \in \mathbb{Z}.$$

Definition 8.8. A module M over a graded field algebra $(V, |0\rangle, Y)$ is called *graded* if there is a diagonalizable operator $H^M \in \text{End } M$ satisfying (cf. (1.6))

$$(8.7) \quad [H^M, Y^M(a, z)] = Y^M(Ha, z) + z\partial_z Y^M(a, z), \quad a \in V.$$

It is called $\mathbb{Z}_+$ -*graded* if all eigenvalues of H^M are non-negative integers.

From now until the end of the section, $(V, |0\rangle, Y)$ will be a $\mathbb{Z}_+$ -graded field algebra and M will be a $\mathbb{Z}_+$ -graded module over it. Then $M = \bigoplus_{k \in \mathbb{Z}_+} M_k$, where $M_k = \{v \in M \mid H^M v = kv\}$ (so $M_k = 0$ for $k \notin \mathbb{Z}_+$). By (1.10), we have:

$$a_n^M M_k \subset M_{k-n}, \quad a \in V, n, k \in \mathbb{Z}_+,$$

where a_n^M is defined similarly to (1.9). In particular,

$$a_0^M v \in M_0, \quad a_n^M v = 0 \quad \text{for all } a \in V, v \in M_0, n > 0.$$

For a homogeneous $a \in V$, $b \in V$, and $r \in \mathbb{Z}$, let us define

$$(8.8) \quad a *_r b = \text{Res}_z z^r (z+1)^{\Delta_a} Y(a, z) b = \sum_{i=0}^{\Delta_a} \binom{\Delta_a}{i} a_{(r+i)} b,$$

and extend it to a bilinear product on V . For $r = -1$, this coincides with Zhu's $*$ -product, and for $r = -2$ with Zhu's $\circ$ -product [Z]. We will also denote $*_{-1}$ as $*$ for short. Using $[H, T] = T$, it is easy to see that

$$(8.9) \quad ((T+H)a) *_r b = -r a *_r b - (r+1) a *_r b, \quad a, b \in V, r \in \mathbb{Z}.$$

In particular, $a *_r b = ((T+H)a) *_r b$.

Proposition 8.9. *Assume that the elements $a, b \in V$, $v \in M_0$ satisfy the associativity relation from Definition 8.1 with $N = \Delta_a$. Then:*

$$(8.10) \quad a_0^M (b_0^M v) = (a *_r b)_0^M v,$$

$$(8.11) \quad 0 = (a *_r b)_0^M v \quad \text{for } r \leq -2.$$

Proof. Equation (8.10) follows immediately from (1.8), (1.9) and (8.5) for $n = \Delta_a - 1$, $N = \Delta_a$, $P = \Delta_b$. Equation (8.11) follows from (8.9), (8.10) and (1.11). $\square$

It follows from Remark 3.12 that the assumption of Proposition 8.9 is satisfied when V is a $\mathbb{Z}_+$ -graded strong field algebra and M is a $\mathbb{Z}_+$ -graded module over it.

Proposition 8.10. *If V is a $\mathbb{Z}_+$ -graded strong field algebra, the space $V *_r V = \text{span}\{a *_r b \mid a, b \in V\}$ is a two-sided ideal with respect to the $*$ -product, and the $*$ -product becomes associative on the quotient.*

The associative algebra $\text{Zh}(V) = V/(V *_r V)$ is called the *Zhu algebra* of V (see [Z]). Then, by Proposition 8.9, M_0 is a module over $\text{Zh}(V)$.

Proof of Proposition 8.10. First of all, notice that by (8.9)

$$(8.12) \quad a *_r b \in V *_r V \quad \text{for all } r \leq -2 \text{ and } a, b \in V.$$

We are going to show that for any $a, b, c \in V$

$$(8.13) \quad (a *_r b) *_s c \equiv a *_r (b *_s c) \pmod{V *_r V} \quad \text{for } r, s \leq -1.$$

We may assume that a and b are homogeneous. Then, using (1.8), we find:

$$\begin{aligned} (a *_r b) *_s c &= \sum_{i=0}^{\Delta_a} \binom{\Delta_a}{i} \text{Res}_u u^s (u+1)^{\Delta_a + \Delta_b - r - i - 1} Y(a_{(r+i)} b, u) c \\ &= \text{Res}_z \text{Res}_u z^r u^s (u+1)^{\Delta_b - r - 1} (z+u+1)^{\Delta_a} Y(Y(a, z) b, u) c. \end{aligned}$$

Now we use (3.4) with $w = -u$. Making the change of variables $x = z + u$, we obtain for any $b' \in V$:

$$\begin{aligned} \text{Res}_z z^r (z + u + 1)^{\Delta_a} i_{z,u} Y(a, z + u) b' &= \text{Res}_x i_{x,u} (x - u)^r (x + 1)^{\Delta_a} Y(a, x) b' \\ &= \sum_{k=0}^{\infty} \binom{r}{k} (-u)^k a *_{r-k} b' \equiv a *_{r-1} b' \pmod{V *_{-2} V} \quad \text{for } r \leq -1. \end{aligned}$$

Therefore, the first term in the right-hand side of (3.4) gives $a *_{r-1} (b *_{-1} c)$ modulo $V *_{-2} V$. The contribution of the second term is zero, because for $r \leq -1$ and $j \geq 0$

$$\text{Res}_z z^r (z + u + 1)^{\Delta_a} \partial_u^j \delta(z - (-u)) \in u^r \mathbb{C}[u^{-1}].$$

This completes the proof of (8.13), and hence of the proposition. $\square$

A large part of Zhu's theory can be generalized from vertex algebras to the case of strong field algebras. For example, we have the following result (cf. [Z]).

Proposition 8.11. *If V is a $\mathbb{Z}_+$ -graded strong field algebra, the functor $M \mapsto M_0$ establishes a one-to-one correspondence between irreducible $\mathbb{Z}_+$ -graded V -modules and irreducible $\text{Zh}(V)$ -modules.*

However, because of Theorem 7.4, the case of strong field algebras is not much different than the case of vertex algebras. Some of the results hold in the more interesting case of field algebras (see e.g. (8.5), (8.6)), but we do not know whether $V *_{-2} V$ is an ideal or $V/(V *_{-2} V)$ is associative in such generality.

Example 8.12. Let V be the tensor product of a $\mathbb{Z}_+$ -graded strong field algebra V' and a unital algebra A (see Example 1.4). Then, although V is not a strong field algebra, Propositions 8.9–8.11 hold for V . One has: $\text{Zh}(V) \simeq \text{Zh}(V') \otimes A$.

Example 8.13. Similarly, let $V = V' \sharp \Gamma$ be the smash product of a $\mathbb{Z}_+$ -graded strong field algebra V' and a group Γ of its automorphisms (see Example 1.5). Then Propositions 8.9–8.11 hold for V . Note that the action of Γ on V' induces an action on its Zhu algebra $\text{Zh}(V')$. One has: $\text{Zh}(V) \simeq \text{Zh}(V') \sharp \Gamma$.

ACKNOWLEDGMENTS

We are grateful to MSRI, where this work was completed, for hospitality. One of the authors wishes to thank V. Schechtman and L. W. Small for very useful discussions and V. L. Popov and E. B. Vinberg for correspondence. We thank A. De Sole for carefully reading the manuscript and suggesting improvements, and C. Dong for pointing out a gap in a previous version of the paper.

REFERENCES

- [B1] R. E. Borcherds, *Vertex algebras, Kac-Moody algebras, and the Monster*, Proc. Nat. Acad. Sci. U.S.A. **83** (1986), no. 10, 3068–3071.
- [B2] R. E. Borcherds, *Quantum vertex algebras*, in “Taniguchi Conference on Mathematics” (Nara, 1998), 51–74, Adv. Stud. Pure Math., **31**, Math. Soc. Japan, Tokyo, 2001.
- [DLM] C. Dong, H. Li, and G. Mason, *Compact automorphism groups of vertex operator algebras*, Internat. Math. Res. Notices **18** (1996), 913–921.
- [DY] C. Dong and G. Yamskulna, *Vertex operator algebras, generalized doubles and dual pairs*, preprint [math.QA/0006005](#), to appear in Math. Z.
- [EK] P. Etingof and D. Kazhdan, *Quantization of Lie bialgebras. V. Quantum vertex operator algebras*, Selecta Math. (N.S.) **6** (2000), 105–130.

- [FR] E. Frenkel and N. Reshetikhin, *Towards deformed chiral algebras*, in Proc. Quantum Group Symp. at XXIth Intern. Coll. on Group Theoretical Methods in Physics, Goslar 1996; `q-alg/9706023`.
- [GMS] V. Gorbounov, F. Malikov, and V. Schechtman, *Gerbes of chiral differential operators. II*, preprint `math.AG/0003170`.
- [K] V. G. Kac, *Vertex algebras for beginners*, AMS University Lecture Series, vol. 10, 1996. Second edition, 1998.
- [KR] V. G. Kac and A. D. Radul, *Representation theory of the vertex algebra $W_{1+\infty}$* , Transform. Groups **1** (1996), 41–70.
- [KT] V. G. Kac and I. T. Todorov, *Affine orbifolds and rational conformal field theory extensions of $W_{1+\infty}$* , Comm. Math. Phys. **190** (1997), 57–111.
- [L] S. Lang, *Algebra*. Third edition, Addison Wesley, Reading, MA, 1993.
- [MS] S. Montgomery and L. W. Small, *Fixed rings of Noetherian rings*, Bull. London Math. Soc. **13** (1981), 33–38.
- [Z] Y. Zhu, *Modular invariance of characters of vertex operator algebras*, J. Amer. Math. Soc. **9** (1996), 237–302.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CALIFORNIA, BERKELEY, CA 94720, USA
E-mail address: `bakalov@math.berkeley.edu`

DEPARTMENT OF MATHEMATICS, MIT, CAMBRIDGE, MA 02139, USA
E-mail address: `kac@math.mit.edu`